

TEFLON CHEMICAL UNSAFE AT SMALLEST DOSES

EPA'S "SAFE" LEVEL
IS HUNDREDS OR
THOUSANDS OF
TIMES TOO WEAK

EWG

AUGUST 2015

By Bill Walker
Investigations Editor

David Andrews
Senior Scientist



www.ewg.org

1436 U Street N.W., Suite 100
Washington, D.C. 20009



Contents

- 3 Report
- 5 PFOA pollution is worldwide – and in people’s blood
- 5 Phased out, but still a threat
- 6 Nationwide water sampling
- 8 Endnotes



www.ewg.org

Designers

Print Layout
Aman Anderson

Web Layout
Taylan Yalniz

About EWG

The Environmental Working Group is the nation’s most effective environmental health research and advocacy organization. Our mission is to conduct original, game-changing research that inspires people, businesses and governments to take action to protect human health and the environment. With your help – and with the help of hundreds of organizations with whom we partner – we are creating a healthier and cleaner environment for the next generation and beyond.

Reprint Permission

To request reprint permission, please email a completed request form to permissionrequests@ewg.org

HEADQUARTERS

1436 U Street N.W., Suite 100
Washington, D.C. 20009
(202) 667-6982

CALIFORNIA OFFICE

2201 Broadway, Suite 308
Oakland, CA 94612

MIDWEST OFFICE

103 E. 6th Street, Suite 201
Ames, IA 50010

SACRAMENTO OFFICE

1107 9th Street, Suite 625
Sacramento, CA 95814

TEFLON CHEMICAL UNSAFE AT SMALLEST DOSES

EPA'S "SAFE" LEVEL IS HUNDREDS OR THOUSANDS OF TIMES TOO WEAK

By Bill Walker, Investigations Editor
David Andrews, Senior Scientist

Every time they drink a glass of tap water, people in the mid-Ohio River Valley of West Virginia and Ohio may be consuming unsafe amounts of an industrial chemical linked to cancer, birth defects, heart disease and other illness. More than a decade after this threat became known, government regulators have failed to set enforceable standards to ensure the water is safe – and now, new science says the danger may be much greater than either residents or regulators thought.

In 2005, DuPont settled a class-action lawsuit brought on behalf of 70,000 mid-Ohio Valley residents for decades of fouling their drinking water with a highly toxic chemical once used to make Teflon. As part of the settlement, DuPont¹ is paying for technology to filter – but not eliminate – the toxin from six area water systems.

Next month, the first of approximately 3,500 personal injury lawsuits² from mid-Ohio Valley residents who got sick from drinking the contaminated water will go to trial. The U.S. Environmental Protection Agency has spent a decade studying the health hazards of the Teflon chemical, known as PFOA,³ but may take another four to six years before even deciding whether to set a legally enforceable maximum pollution level for drinking water. In 2009, EPA set a non-enforceable Provisional Health Advisory level – a temporary, voluntary standard to help utilities and health officials decide when to take action to reduce peoples' exposure – but the agency didn't follow up with a rule that carries the force of law. That advisory level remains the only federal guidance on how much PFOA is safe in drinking water.

Now two leading environmental health scientists have published research with alarming implications: PFOA contamination of drinking water is a much more serious threat to health, both in the mid-Ohio Valley and nationwide, than previously thought. Their research finds that even very tiny concentrations of PFOA – below the reporting limit required by EPA's tests of public water supplies – are harmful. This means that EPA's health advisory level is hundreds or thousands of times too weak to fully protect human health with an adequate margin of safety. The implications:

- The settlement set a "trigger level" of PFOA to determine which West Virginia and Ohio water systems DuPont would pay to have filtered. That trigger level is more than 160 times the amount the new research says is safe. Filtration has cut PFOA to almost non-detectable levels in many of the water systems covered by the settlement, but even the least contamination measured exceeds the amount the new research says is safe.
- Only those people in the mid-Ohio Valley who became sick after drinking water at or above that trigger level are eligible to be plaintiffs in the upcoming trials, but the new science indicates that many more people who drank less heavily contaminated water could also have been harmed. (They may file suits in the future, but will not have the benefit of the extraordinary concession DuPont has made for the upcoming trials – that PFOA at the trigger level can cause certain diseases.)

- Two water systems in the region – Parkersburg and Vienna, W. Va. – were not covered by the 2005 settlement because their PFOA contamination at that time was just under the trigger level. Without benefit of the state-of-the-art filtration technology installed in other systems, water samples taken in Vienna in May 2007 had PFOA levels more than 180 times higher than the new research says is safe,⁴ and those taken in Parkersburg last September were more than 130 times the new safe level.
- Since 2013, an EPA testing program has found [PFOA in 94 public water systems in 27 states](#). These systems provide drinking water to more than 6.5 million people. The vast majority of water samples tested had no detectable level of PFOA, and in every system with PFOA the level was well below the EPA advisory level. But among the samples with PFOA, statewide average levels ranged between five times and 175 times the level described by the new research as safe.⁵
- To replace the provisional health advisory limit set in 2009, EPA officials are in the process of establishing a long-term health advisory level for PFOA in drinking water. A draft EPA study released last year suggested that the advisory level being considered could be more than 300 times the new safe limit. Whatever the new advisory level turns out to be, it will remain voluntary: agency officials have said they could take until 2021 years to decide whether to attempt to set a legally enforceable maximum for PFOA.⁶

PFOA POLLUTION IS WORLDWIDE – AND IN PEOPLE’S BLOOD

Through Teflon’s use in hundreds of household products – carpets, clothing, food wrappers and many more – PFOA and closely related chemicals have spread to the remote corners of Earth, contaminating

TABLE 1.

Comparison of various PFOA levels with the safe level suggested by new research

	Level of PFOA (parts per billion)	Level of PFOA compared to safe level in new research
EPA Provisional (temporary) Health Advisory (2009)	0.4 ppb	1,333X
EPA Draft of Chronic (long-term) Health Advisory (2014)	0.1 ppb	333X
“Trigger level” for filtration of mid-Ohio Valley water systems covered by 2006 settlement	0.05 ppb	166X
Threshold for drinking water consumption by plaintiffs in upcoming personal injury trials	0.05 ppb	166x
PFOA level in Parkersburg, W.Va. water (2014)	0.0412 ppb	137X
PFOA level in Vienna, W.Va. water (2007)	0.056 ppb	187X
Lowest average level of detection of PFOA in state water systems, 2013-2015 (Va.)	0.0109 ppb	5x
Highest average level of detection of PFOA in state water systems, 2013-2014 (W.Va.)	0.0383 ppb	174X

Sources: EWG, from Grandjean and Clapp 2015; EPA 2009, 2015; Bilott to EPA 2015.

the blood of virtually all Americans and even passing through the umbilical cord to unborn babies in the womb.

PFOA has been linked to kidney and testicular cancers, birth defects, damage to the immune system, heart and thyroid disease, complications during pregnancy and other serious illnesses and conditions. It is hazardous at tiny doses: EPA's health advisory level for drinking water is 0.4 parts per billion. (A part per billion, or ppb, is less than a teaspoon in an Olympic-sized swimming pool.)

Last June, the scientific journal *New Solutions* published a paper by Philippe Grandjean of the Harvard School of Public Health and Richard Clapp of the University of Massachusetts-Lowell, reviewing the research EPA used to set its health advisory level and comparing it to more recent studies.⁷

In setting its advisory level, EPA relied studies before 2008 on the effects of PFOA and the closely related compound PFOS⁸, formerly used in 3M's Scotchgard, on the weight of the liver and kidneys of rats and mice. PFOA toxicity testing has often been done using rats but female rats eliminate PFOA from their bodies much faster than people, so rats are not an ideal species for studying human developmental effects.⁹ EPA also considered PFOA only "suggestive" of carcinogenicity, but both an external review panel appointed by the agency and a science panel funded by the DuPont settlement¹⁰ later declared that PFOA is a "likely" cause of cancer.

Grandjean and Clapp cited a newer study by the National Toxicology Program, EPA and the University of North Carolina. This research concluded that PFOA could disrupt hormones and suggested a possible link to breast cancer.¹¹ That study dosed mice in the womb with very low levels of PFOA during critical development periods and could not find a level so low it did not cause harm. The scientists also drew on Grandjean's 2013 study of more than 400 children in the Faeroe Islands of the North Atlantic. These children's diet was heavy in PFOA-contaminated fish. The results suggested that PFOA exposure could reduce the effectiveness of childhood vaccines.¹²

In another sign of the growing scientific recognition that PFOA is more harmful than previously thought, the National Toxicology Program recently announced a [systematic re-evaluation](#) of the chemical's effect on

the immune system. The program's Office of Health Assessment and Translation issued a call for the submission of ongoing or upcoming studies to be considered in the evaluation and for the nomination of scientists for an expert panel to review the findings.

Grandjean and Clapp suggested that the EPA's approach in 2009 led to a presumed safe level "at least two orders of magnitude" higher than the newer studies indicate would protect human health with an adequate margin of safety. Grandjean and Clapp termed 0.001 ppb the "approximate" safe level for PFOA, but EWG's calculations from their data yielded a level of 0.0003 ppb – lower than the EPA advisory level by a factor of more than 1,300.

PHASED OUT, BUT STILL A THREAT

Both PFOA and PFOS belong to a class of non-stick, waterproof, grease-proof chemicals historically called PFCs.¹³ Under pressure from EPA, 3M stopped making PFOS in 2002, and in 2005 DuPont agreed to phase out PFOA by this year. Those two chemicals are no longer produced in the U.S.

But DuPont and other chemical companies are marketing a new generation of PFCs with similar chemical structures. The few studies conducted on these new chemicals show that they may also have serious health risks. But the weak and outdated federal Toxic Substances Control Act has allowed them onto the market without adequate safety testing. Grandjean and Clapp wrote that "the greatly underestimated health risks from [PFOA] and [PFOS] illustrate the public health implications of assuming the safety of incompletely tested industrial chemicals."

The new science demands urgent action to set stricter and legally enforceable limits on PFOA in drinking water. The threat to public health is most severe in the mid-Ohio Valley and the state-level test results reveal a problem nationwide. But federal regulators are moving at a glacial pace.

EPA was first alerted to PFOA pollution in the mid-Ohio Valley in 2001. Not until 2009 did it produce the current advisory level, which it called a "reasonable, health-based hazard concentration above which

action should be taken to reduce exposure.”¹⁴ Last year, EPA released a draft¹⁵ of its proposed “reference dose” – an estimate of how much a person can safely consume daily over a lifetime. That proposed reference dose would translate to a legal limit for PFOA of 0.1 ppb.

That’s a quarter of the current advisory level but still more than 300 times the safe level put forth by Grandjean and Clapp. What’s more, the EPA draft study says that PFOA exposure is only “suggestive of carcinogenicity,” again disregarding the findings in 2006 of EPA’s own Science Advisory Panel that PFOA is a “likely” human carcinogen.

EPA’s draft study and the findings of the nationwide water sampling will drive the decision of whether to set a legal limit for PFOA in drinking water, and it is far from certain that the agency will act. In February of last year, Nancy Stoner, EPA’s acting water chief, told *Inside EPA*: “The agency expects that we will have sufficient information to determine whether it is appropriate to develop a drinking water regulation for PFOA within the next 5 to 7 years.”¹⁶

In other words, EPA officials think they need as much as seven years to decide whether to even propose a legal limit for PFOA. If they do make a proposal, the rule-making process is so protracted that many more years could elapse without a legal maximum for a compound whose threat to health at low doses has been confirmed by scores of peer-reviewed studies.

NATIONWIDE WATER SAMPLING

Nationwide sampling for PFOA, PFOS and four other PFCs¹⁷ in drinking water began in 2013, under an EPA program¹⁸ that periodically requires all U.S. public water systems serving 10,000 or more people to test for contaminants that are not yet regulated. Through July of this year, the current round of the program had tested more than 29,000 samples. Fewer than one percent of the samples had detectable levels of PFOA. But critics, including the New Jersey Department of Environmental Protection and the American Water Works Association, said in

comments to EPA that the tests were not designed to detect very low levels of the chemicals.¹⁹

In 2006 New Jersey conducted its own tests for PFOA and other PFCs in drinking water. The methods used by New Jersey officials were approximately 10 times the sensitivity of those specified by EPA. The less sensitive EPA tests and reporting threshold used for the nationwide sampling program would have missed almost three-fourths of the PFC water contamination New Jersey officials found in their state. New Jersey’s testing prompted state regulators to set their own non-enforceable health advisory level for PFOA in drinking water of 0.04 ppb – ten times more protective than EPA’s advisory level, but still more than 130 times the amount Grandjean and Clapp said is safe.

The results of the nationwide tests could mean that EPA will decline to propose an enforceable rule for PFOA in drinking water, because the federal Safe Drinking Water Act requires that when the agency decides whether to issue a rule with the force of law, it must consider “the frequency and level of contaminant occurrence in public drinking water systems.”²⁰ Yet the new research shows that the average of levels detected in each state are between five and 175 times too high to be considered safe.

The EPA-mandated tests by local water systems found PFOA in 27 states, in 94 water systems serving 6.5 million Americans. The results show that outside of the mid-Ohio Valley, New Jersey and California have the most widespread PFOA contamination. Testing in both states found PFOA in 14 water systems, serving more than 1.4 million people in California and more than 1.3 million people in New Jersey. By the most conservative estimate, the average level of PFOA detected in California samples was 14 times the level Grandjean and Clapp say is safe, and in New Jersey it was 30 times the new safe level.

The table below shows average levels of PFOA in water samples, and by how much they exceed the new safe level according to Grandjean and Clapp. [EWG has also produced an interactive map that shows nationwide detections of PFOA, PFOS and four other PFCs.](#)

TABLE 2

Nationwide sampling found that statewide average detections of PFOA in drinking water greatly exceed the safe level established by the new research.

State	Number of systems with water samples with PFOA	Number of water samples with PFOA	Percent of samples with PFOA detected	Population served by drinking water systems with PFOA ^a	Range of average level of PFOA in samples (parts per billion) ^b	Average level of PFOA compared to safe level in new research
Alabama	12	26	25%	312,522	0.0081-0.0231	27-77X
Arizona	4	14	15%	217,218	0.006-0.0219	20-73X
California	14	39	14%	1,441,298	0.0041-0.035	14-117X
Colorado	3	52	78%	57,343	0.0361-0.0397	120-132X
Delaware	2	12	30%	320,494	0.0172-0.0269	57-90X
Florida	2	4	7%	371,913	0.0028-0.017	9-57X
Georgia	2	3	100%	94,674	0.0367	122X
Illinois	3	3	33%	135,753	0.0136-0.0209	45-70X
Kentucky	1	2	25%	730,611	0.005-0.0131	17-44X
Massachusetts	3	7	21%	103,762	0.0062-0.0168	21-56X
Maryland	1	1	10%	104,567	0.0021-0.012	7-40X
Minnesota	4	6	26%	143,637	0.0256-0.0346	85-115X
North Carolina	5	8	9%	225,262	0.002-0.0192	7-64X
New Hampshire	2	2	17%	53,000	0.0091-0.0184	30-61X
New Jersey	14	60	33%	1,334,413	0.0091-0.0272	30-91X
New York	3	4	8%	174,000	0.0028-0.0163	9-54X
Ohio	3	3	33%	79,337	0.008-0.0153	27-51X
Oklahoma	1	1	33%	20,307	0.0113-0.0182	38-61X
Pennsylvania	5	24	25%	221,121	0.0193-0.0337	64-112X
Rhode Island	1	1	25%	21,900	0.0203-0.0281	68-94X
South Carolina	1	1	25%	24,904	0.006-0.0138	20-46X
Tennessee	1	1	25%	139,110	0.005-0.0128	17-43X
Texas	1	1	25%	11,489	0.0066-0.0144	22-48X
Virginia	1	1	7%	47,574	0.0016-0.0122	5-41X
Washington	3	6	23%	109,527	0.0067-0.0163	22-54X
Wisconsin	1	3	21%	30,100	0.0063-0.0153	21-51X
West Virginia	1	2	100%	34,251	0.0522	174X
Total	94	287	23%	6,560,087		

NOTES

a - From the EPA Safe Drinking Water Information System 2013 Inventory.

b - EPA only required reporting of PFOA levels above 0.02 ppb, even if the laboratories testing the samples had lower limits of detection. The average PFOA values for each state are provided as a range, calculated using zero and 0.02 ppb as the value for all non-detects.

ENDNOTES

1. In July 2015 DuPont spun off its fluorochemicals business to a new company named Chemours, which will also inherit DuPont's liabilities for PFOA pollution.
2. The thousands of individual suits have been consolidated into one case in federal district court in Columbus, Ohio. Two individual cases will be tried first, starting Sept. 14.
3. Perfluorooctanoic acid. DuPont referred to the compound as C8, and much of the regulatory and scientific literature on the pollution in the mid-Ohio Valley uses that designation.
4. Letter from attorney Robert A. Bilott to EPA regional administrators and the West Virginia Department of Environmental Protection, January 20, 2015.
5. U.S. EPA, Unregulated Contaminants Monitoring Rule 3 (2013-2015) Occurrence Data. June 2015. Available: water.epa.gov/lawsregs/rulesregs/sdwa/ucmr/data.cfm
6. Suzanne Yohannan, EPA Proposes First-Time Risk Values for Chronic PFC Exposures. InsideEPA.com, February 28, 2014. Available: insideepa.com/daily-news/epa-proposes-first-time-risk-values-chronic-pfc-exposures.
7. Philippe Grandjean and Richard Clapp, Perfluorinated Alkyl Substances: Emerging Insights Into Health Risks. *New Solutions: A Journal of Environmental and Occupational Health Policy*, June 17, 2015. Available: new.sagepub.com/content/25/2/147
8. Perfluorooctanesulfonic acid.
9. Post GB, Cohn PD, Copper KR. Perfluorooctanoic acid (PFOA), an emerging drinking water contaminant: a critical review of recent literature. *Environmental Research*, July 2012. Available: <http://www.ncbi.nlm.nih.gov/pubmed/22560884>
10. The C8 Science Panel, three independent experts who for seven years studied the chemical's effects on 70,000 residents of the mid-Ohio Valley.
11. D.K. Tucker et al, The mammary gland is a sensitive pubertal target in CD-1 and C57Bl/6 mice following perinatal perfluorooctanoic acid (PFOA) exposure. *Reproductive Toxicology*, July 2015. Available: www.ncbi.nlm.nih.gov/pubmed/25499722
12. Philippe Grandjean and Esben Budtz-Jørgensen, Immunotoxicity of perfluorinated alkylates: calculation of benchmark doses based on serum concentrations in children. *Environmental Health*, April 19, 2013. Available: www.ehjournal.net/content/12/1/35
13. Polyfluorinated chemicals. Most researchers now use the broader term poly- and perfluoroalkyl substances, or PFAS.
14. U.S. EPA, Provisional Health Advisories for PFOA and PFOS. January 8, 2009. Available: water.epa.gov/action/advisories/drinking/upload/2009_01_15_criteria_drinking_pha-PFOA_PFOS.pdf
15. U.S. EPA, Health Effects Document for Perfluorooctanoic Acid (PFOA), draft, February 2014. Available: [peerreview.versar.com/epa/pfoa/pdf/Health-Effects-Document-for-Perfluorooctanoic-Acid-\(PFOA\).pdf](http://peerreview.versar.com/epa/pfoa/pdf/Health-Effects-Document-for-Perfluorooctanoic-Acid-(PFOA).pdf)
16. Yohannan, InsideEPA.com, op. cit. Available: insideepa.com/daily-news/epa-proposes-first-time-risk-values-chronic-pfc-exposures.
17. PFBS, PFHpA, PFHxS and PFNA. PFOS, the only other PFC for which EPA has set a Provisional Health Advisory level – 0.2 ppb, half that of PFOA – was detected in 234 samples from 87 water systems.
18. The Unregulated Contaminant Monitoring Rule.
19. U.S. EPA, Revisions to the Unregulated Contaminant Monitoring Regulations (UCMR) for Public Water Systems. Comments from American Water Works Association, Fluorocouncil and New Jersey Department of Environmental Protection. Docket ID: EPA-HQ-OW-2009-0090.
20. U.S. EPA, Regulating Public Water Systems and Contaminants Under the Safe Drinking Water Act. Available: water.epa.gov/lawsregs/rulesregs/regulatingcontaminants/basicinformation.cfm